



## Tammar Wallaby *Macropus eugenii* (Desmarest, 1817)



Photo: Babs & Bert Wells/DEC

### Size

#### Kangaroo Island wallabies

Head and body length

590-680 mm in males

520-630 mm in females

Tail length

380-450 mm in males

330-440 mm in females

### Weight

#### Western Australian wallabies

2.9-6.1 kg in males

2.3-4.3 kg in females

### Subspecies

Three subspecies are recognised:

- *Macropus eugenii derbianus* (Western Australia)
- *Macropus eugenii decres* (Kangaroo Island, South Australia)
- *Macropus eugenii eugenii* (South Australian mainland)

### Description

Dark, grizzled grey-brown above, becoming rufous on the sides of the body and the limbs, especially in males. Pale grey-buff below.

### Other Common Names

Dama Wallaby (South Australia)

### Distribution

The Western Australian subspecies of the Tammar Wallaby was previously distributed throughout most of the south-west of Western Australia from Kalbarri National Park to Cape Arid on the south coast and extending to western parts of the Wheat belt.

The Tammar Wallaby is currently known to inhabit three islands in the Houtman Abrolhos group (East and West Wallabi Island, and an introduced population on North Island), Garden Island near Perth, Middle and North Twin Peak Islands in the Archipelago of the Recherche, and several sites on the mainland - including, Dryandra, Boyagin, Tutanning, Batalling (reintroduced), Perup, private property near Pingelly, Jaloran Road timber reserve near Wagin, Hopetoun, Stirling Range National Park, and Fitzgerald River National Park. The Tammar Wallaby remains relatively abundant at these sites which are subject to fox control.

They have been reintroduced to the Darling scarp near Dwellingup, Julimar Forest near Bindoon, state forest east of Manjimup, Avon Valley National Park, Walyunga National Park, Nambung National Park and to Karakamia and Paruna Sanctuaries.

For further information regarding the distribution of this species please refer to [www.naturemap.dpaw.wa.gov.au](http://www.naturemap.dpaw.wa.gov.au)

### Habitat

Dense, low vegetation for daytime shelter and open grassy areas for feeding. This species inhabits coastal scrub, heath, dry sclerophyll forest and thickets in mallee and woodland.

### Behaviour

The Tammar Wallaby is nocturnal. During the day they rest in low scrub and, although they begin to move at dusk, they do not leave the scrub until after dark and return to it before dawn. Individuals

have defined home ranges (area of occupancy) that overlap the home ranges of others. While several wallabies have been observed feeding in the same area, no social grouping has been observed except between females and their young at foot.

## Diet

The Tammar Wallaby is herbivorous, and its diet consists mostly of grasses. Evidence suggests it requires drinking water for its survival. One study found up to 24 plant species in the diet, including *Gastrolobium bilobum*, *Danthonia setacea*, and *Corymbia calophylla*.

## Breeding

The breeding cycle is well known only for the Kangaroo Island subspecies (*Macropus eugenii. decres*). The Tammar Wallaby is one of only two macropod species that show a strictly seasonal pattern of breeding. Most young are born from late January to March and within a few hours of giving birth the female mates and the resulting embryo remains dormant during lactation. The dormant embryos are reactivated within a few days after mid December, and the young enter the pouch about 40 days later, 12 months after the mating at which they were conceived. The single young is suckled in the pouch for eight to nine months and leaves the pouch in September or October (October – November in Western Australia). Females become mature at about nine months while they are still suckling, but males do not become mature until nearly two years old. The rate of reproduction is high with more than 90% of all females carrying a pouch young by the end of the breeding season. In some years many pouch young are lost, particularly by one-year-old females. In all years mortality is high among juveniles during their first summer and may reach 40%.

## Threatening processes

Fox predation, and loss of suitable thickets due to inappropriate fire regimes and land clearing.

## Conservation status

- *Macropus eugenii*

IUCN Red List of Threatened Species

Least concern

- *Macropus eugenii derbianus*

Not Listed

- *Macropus eugenii eugenii*

Environment Protection and Biodiversity Conservation Act 1999

Extinct

(This subspecies has been recently reintroduced to South Australia from an introduced population in New Zealand)

- *Macropus eugenii decres*

Not Listed

## Management

The following management actions are recommended:

- Monitor Tammar Wallaby populations at forest and woodland sites as part of threatened species reintroduction programs and broad scale fox control programs.
- Implement appropriate fire regimes to ensure regeneration of shelter thickets.

## Other interesting facts

- The Tammar Wallaby was the first "kangaroo" seen by Europeans (in 1629) by the crew of the Dutch ship *Batavia* stranded near the Wallabi Islands in the Houtman Abrolhos off Geraldton.
- Tammar Wallabies were so named because they were once common in Tamma (*Allocasuarina campestris*) thickets.
- Lifespan is approximately 11 years for males and 14 years for females.
- In semi arid areas such as the Abrolhos Islands, Tammar Wallabies are able to drink sea water when fresh water is unavailable.

## Selected references

Anon. (1998). Bush Telegraph: Two More Mammals off the Threatened List. *Landscape* 13(4): 5.

Hinds, L.A. (2008). Tammar Wallaby. In Van Dyck, S. and R. Strahan (Eds.) *The Mammals of Australia*. Reed New Holland. Sydney

Maxwell S., Burbidge A.A, Morris K. (1996). *The 1996 Action Plan for Australian Marsupials and Monotremes*. Wildlife Australia, Canberra.

Poole, W. E., Wood, J. T. and Sims, N. G. (1991). Distribution of the Tammar and relationships of populations as determined by cranial morphometrics. *Wildlife Research* 18, 625-639.

Shephard, K.A., Wardell-Johnson, G.W., Longeragan, W.A. and Bell, D.T. (1997). Diet of herbivorous marsupials in a *Eucalyptus marginata* forest and their impact on the understorey vegetation. *Journal of the Royal Society of Western Australia* 80: 47-54.

Last updated 8 February 2012, for further enquiries please contact [fauna@dpaw.wa.gov.au](mailto:fauna@dpaw.wa.gov.au)

## Further information

Contact your local office of the Department of Environment and Conservation.

See the department's website for the latest information: [www.dec.wa.gov.au](http://www.dec.wa.gov.au).

*Disclaimer: This publication may be of assistance to you but the Government of Western Australia and its officers do not guarantee that the publication is without flaw of any kind or is wholly appropriate for your particular purposes and therefore disclaims all liability for any error, loss or other consequence which may arise from you relying on any information in this publication.*



Department of  
Environment and Conservation

